Comparison of different commercial fruit beverage characteristics in Iran

Mahkameh Yousef lavi¹, Leila tabatabaei moradi,*¹ Morvarid Yousef¹, Leila Nateghi¹ ¹Department of Food Science and Technology Varamin-Pishva Branch, Islamic Azad University Varamin, Iran e-mail : Mah_lavi@yahoo.com , Leila94132@yahoo.com , Yousefi.morvarid@ yahoo.com, leylanateghi@yahoo.com

Abstract

In this study, it was aimed to measure the physic- chemical characteristics of six kind of Iranian commercial fruit beverage in clouded cherry, apple, pineapple, mango, orange and grape that were obtained from local store. In this samples mean value for pH, total sugar, fat, calcium and phosphorus, total solid and total soluble solid content was 3.32, 8.26%,0, 0.03%,0.05%14.5 %,and16.40 % respectively.

Key words: fruit beverage, chemical properties.

Introductions

Fruit juice is a popular beverage in the world. Generally in juice processing, initially fruits are milled which is subsequently extracted or pressed and a cloudy liquid obtained. This liquid is clarified, screened and then pasteurized.(Zarate-rodriguez and Ortega-Rivas 1999). Historically, the conniption of fruit juices began with usage of orange juice, due to prevent scurvy, as a source of vitamin ascorbic acid. Today markets are flooded with different kind of juices apple, guava, mango, litchi, pineapple, grape, etc. The key reason for increased usage is raising level of health awareness and changing people lifestyles .They believes that these products provide valuable nutrition. Child preference, convenience, easy availability, and naturalness that have given fruit juice and drink industry a booming growth. (Gupta and Gupta, 2008). According to Codex Alimentations Commission, Fruit juice is the unfermented beverage but fermentable product that produced from the edible part of sound, fresh fruit and appropriately matures. Also some product may be processed with seeds, pips and peel, which is not generally incorporated in the product, but some parts of seeds, pips and peel, which cannot be omitted by Good Manufacturing Practices will be acceptable. The beverage is produced by suitable processes, which preserve the essential nutritional, chemical, physical and organoleptical characteristics of the juices of the vegetable and fruit from which it comes. The juice may be clear or cloudy and may have restored volatile flavour and aromatic substances, all of them must be prepared from the same kind of fruit. Fruit juice was produces in two ways the first directly expressed by mechanical extraction processes, and the second, Fruit juice from concentrate that made by mixing concentrated fruit juice with water, also Fruit purée may be used in the preparing of fruit Juices .it is the unfermented but fermentable food stuff obtained by processes include grinding, sieving, milling the edible part of the fruit without eliminating the juice. The fruit juices shall have the characteristic aroma, color and flavor of juice from the same fruit from that it is made. (Codex standard, 2005). According to American Academy of Pediatrics (AAP) fruit juice defines as natural or 100% concentrate without addition of sweating agent. Anything less than 100% concentrate is known as beverage, cocktail or drink (AAP, 2001). Fruit beverages are defined as sweetened calorically drink with a low content of fruit juice or juice flavoring with carbonated water (Pompkin et al ,2006). Fruit drinks have less than 20% concentrate and may be fortified with nutrient such as calcium or vitamin C.Fruit beverage, even 100% juice is not equivalent to natyral and whole fruits because Fruits supply photochemical and fibers to diet, that are removed in juices. Fruit drinks are thought to be valuable and suitable source of minerals .If fruit drinks consumed by children in optimum level, can be a important part of balanced diet . different Studies have shown that presence of flavinoids and vitamin C in juices have beneficial health effects such as decreasing the risk of heart disease and cancer (Ames .1998). Vitamin C by increasing absorption of iron to about double can reduce the anemia in population with low iron content in its diet (Hollman et al ,1996).. This product is perceived as a nutritious, all day beverage, wholesome and is gaining in popularity and use with high increase in sales in recent years. Therefore the aim of this was compare of the physicochemical characteristics properties of some different commercially beverage in Iran.

Material and Methods:

Material

Randomly selected different stores located in Tehran and different kind of fruit beverage include apple, pineapple, mango ,peach ,cherry and orange were obtained from local store. Sampling of each product was down according to International standard number 326.

Methods:

The pH value of samples was measured using pH meter .PH , titratable acidity ,moisture , protein and ash was measured according to the method of international standard . national standars number for grape , apple , pineapple , mango,cherry and orange is1624, 365,10241,10554, 6801 and 507.Calcium and phosphors was measured according to AOAC, 1990.

Data Analysis

Data collected from the aforementioned study samples were analyzed based on 0.05% coefficient of error by a software program. The data analysis was performed using MINITAB statistical software, release 14.2 (MINITAB Inc., state college, PA and USA). At first such software program proved samples normal conditions and then the significant difference among data was precisely studied via Anova – one – way test and p-value was determined.

Results and Discussion

Table 1 and 2 shows the levels of physic -chemical properties of different commercially beverage include cherry, mango, Pineapple, orange, grape and apple. Acidity is a key factor of beverage and juices, and is also easily determined. Acidity of the beverage has an influence on taste and mouth feel of the product. Very high acidity may render the product unpleasant and too sharp and on the other hand if the acidity is very low, the beverage will lack freshness. Between the samples cherry drink show the highest acidity content (1.21%) and statistical measurement showed significant deference between cherry with other samples ($p \le 0.05$). Acceptable limit in international standard for acidity has shown in table 1 and according to this standard all of the samples were acceptable. Pilo et al in 2009 repoted that the acidity of the orange and grape juice varied from 0.40 to 1.40%, and from 0.40 to 1.02 %, respectively, that was agreed with this result. The different kind of acidity that governs this protection is the pH that shows the level of H+ ions in a product. The pH is referring to the different reactions that occur in the solution. These two factors are different: the pH define as inverse logarithmic scale it means if pH is reduced by 1 part, the H+ concentration is multiplied by 10 (Jolicoeur, 1997). Fore the samples the min value of pH was 3.32 and statistically more sour taste than other beverage. Also according to standard all of the samples were suitable. The beverage in this study showed pH values similar to those found by other studied. Pilo et al in 2009 found that The pH of the orange and grape juice varied from 2.94 to 3.85, and from 2.85 to 3.95, respectively. Adou etall ,2012 reported that pH and acidity of apple juice was 4.4 , 0.6 . Also Zarate-Rodroguez et al ,1999 and Carvlho et al , 2011 both of them reported that acidity of apple juice was0.5 % . El-Faki and Eisa, 2010, approved that pH of cherry was about 3% that was agreed with our study. Akhter et al, 2010, found that acidity of mango juice was about 0.15 % that was lower than this result. The pH and acidity of pineapple beverage of this study was similar to the results found by Lee, 2005 that was 0.47%, 3.75 respectively. Generally beverage are fat free or has low level of fat, in this study all of samples were far free. The average level of protein content was 0.33 % and it varies between 0.23 5 to 0.50 5 and statistically were different ($p \le 0/05$). The protein content of beverage was low; of course, this product isn't main source of protein. Total solid (TS) define as weight of whole solids component include organic or inorganic and suspended or dissolved per unit volume of water that generally measured by the evaporation of a determined volume of water at 105 C up to reached the constant weight .TS content in different beverage range from 10.30 5 to 17.32% that refer to Pineapple and mango respectively, it means mango beverage was more concentrated than another samples .Comparison of this parameter shows a significant variation (P<0.05) between the samples . According to inter national standard the lowest level for TS for cherry, pineapple and apple is 12.5%, and for orange and grape is 10.5 % and respectively, therefore all of samples except 15.5% pineapple sample was according to inter national standard. Akhter et al, 2010 ,reported that Ts of mango juice was about 15 %. Adou et al ,2012 aproved that TS of apple juice was about 9 % that was lower than this study.Lee, 2005 has observed that TS content of pineapple juice was 13.64 % that was little higher than our result .Refract index directly show transparency and directly refer to total solid. Refract index of all samples statistically were similar and its ranged was 1.352 (pineapple)and 1.3653(cherry).Costescu et al ,2006 reported that Refract index of orange juice was1.347, that was similar with our study. Total sugar content of the entire sample statistically was similar $(p \le 0/05)$, and its min value was 8.26 %. Total sugar for pineapple and mango was 7.31 5 and 9.42 % respectively. According to the international standard minimum percent of total sugar for pineapple and mango is 12 % and 11 5 respectively and therefore both of them not acceptable, but another samples were in acceptable range. Akhter et al, 2010 ,found that total sugar of mango juice was about 12

% that was higher than our result.Higher content sugar

redoctant related to mango beverage and this difference with

all of them were similar (p≤0.05), However the lowest pH

refer to cherry that was drink 2.98 It means cherry drink has

other samples was significant (p≤0/05). According to international standard at least level for sugar redoctant for mango and grape juice are 3.5 % and 13 % respectively and so both of this samples not acceptable .Akhter et al, 2010 reported that redoctant sugar of mango juice was 2.32 % that was very lower than our result .Carvlho et al, 2011, found that redoctant sugar of apple juice was 12.5 % that was higher than this result .In relation to soluble solids (Brix), a relative wide range of variation was found and statistically analysis shoe significant difference (P<0.05). The highest Brix refer to cherry that was 19.54 % and the lowest related to mango beverage that was 12.40 %. The high content of total soluble solids found in cherry is possibly due to the addition of high level sugar in excess to the beverage, because cherry in nature has sour taste. According to international standard the lowest acceptable Brix for mango is 13.5 %, in term of Brix, mango beverage not acceptable, but other samples are in the acceptable range. Akhter et al, 2010, reported that Brix of mango juice was t 15.5 % that was higher than our study. Pilo et al in 2009 found that the Brix of orange and grape juice varied from 4.93 to 17.45 and from 9.8 to 17.06°, respectively. Adou et al, 2012 and Zarate-Rodrigues et al, 1999 reported that Brix of apple juice was 10 % and 12 % respectively, El-Faki and Eisa, 2010, approved that Brix of cherry was 13%, Also Lee, 2005 found that Brix of pineapple juice was 9 % that all of them were a litter lower than this study .Mango beverage has significant higher ash content than other samples ($p \le 0/05$) and it was 0.81%. According to national standard all of samples was in acceptable range. Adou et al ,2012 reported that ash content of apples juice sample was 1.5%, that was very higher than this study and also higher than international standard .Calcium is an important mineral in the human life, necessary for suitable growth and firmness of the teeth and skeleton as well as muscle and nerve.. The min value of Calcium and Phosphorous was 0.03 % and 0.05 % receptively and all of them were similar.

Conclusions

This study showed the physico-chemical properties of commercial fruit beverages in Iran.

This product has nutritional potential of this fruit such as ash, organic acids, sugar and dry matter. Juices showed a significant difference in all measured factors that related to natural of fruit

Reference

Adou, M., Tetchi , F.A., Gbane , T., Kouassi, K.N., Amani, N.G (2012) Physic – chemical characterization oc cashew apple juice (ANACARDIUM OCCIDENTALE, L.) from yamoussoukro (CÔTE D'IVOIRE) Innovative Romanian Food Biotechnology, 11, 32-43.

Akhter,S., Abid, H., Yasmin,A., Masood,S.(2010) Preparation and evaluation of physical and chemical characteristics of instant mango juice powder. Pak. J. Biochem. Mol. Biol. 2010; 43(2): 58-60

American Academy of Pediatrics. The use and misuses of fruit juices in Pediatrics. Pediatrics 2001; 107: 1210-1213.

Ames BN.(1998) Micronutrients prevent cancer and delay aging. Toxicol Lett; 102-103: 5-18.

Carvlho,C.V.,Chiquetto,N.C.(2011)Foresight of physicalchemical characteristics of apple juiceblends appointed to sparkling drink elaboration . Ciênc. T 188 ecnol. Aliment., Campinas, 31,1, 188-193.

Costescu, C., Pârvu , D., Riviş , A(2006) the determination of some physical – chemical characteristics fore orange , grape fruit and tomato juice .Journal of Agroalimentary Processes and Technologies, 3 , 2 , 429-432.

Codex standard (2005). Codex general standard for fruit juice and nectars, number, 247.

Gupta, H and Gupta, P. (2008) Fruit Drinks: How Healthy and Safe? INDIAN PEDIATRICS ,45 ,17 , 214 -217.

El-Faki A. E., Eisa, E. S.(2010) Physico-Chemical Characteristics of Some Soft Drinks of Sudan During Shelf Life. Journal of Science and Technology 11, 2, 57-62.

Hollman PC, Hertlog MG, Kalan MB.(1996) Role of dietary flavinoids in protection against cancer and coronary heart disease. Biochem Soc Trans , 24: 785-789.

Jolicoeur, C. (1997) Acidity and pH of apple juice. cjoli@gmc.ulaval.ca <u>http://cjoliprsf.awardspace.biz</u>.

Lee S.Y.(2005) Physical and chemical properties of pineapple juice, agglomerate and drink. Malaysian Agricultural Research and Development Institute .

Pompkin BM, Armstrong LE, Bray GM, Caballero B, Frei B, Willett WC.(2006) A new proposed guidance system for beverage consumption in the United States. Am J Clin; 83: 529-542.

Zarate-Rodriguez ,E.,Ortega-Rivas ,E. (1999) Quality changes in apple jouice as related to nonthermal processing .

Table 1: p	hysicochemical	properties o	of fruit beverage
	2		0

Properties (%)	cherry	Cherry	mango	Mango	Pineapple	pineapple
		standard		standard		standard
Total solid	14.00 ± 2.42^{b}	Min 12.5	17.32 ± 1.40^{a}	-	10.30 ± 3.12^{b}	Min 12.5
pН	$2.98{\pm}0.01^{a}$	2.6 - 3.7	3.31 ± 0.00^{a}	Max 3.7	3.45 ± 0.12^{a}	3-4
Refract index	1.3653 ± 0.00^{a}		1.358 ± 0.01^{a}		1.352 ± 0.00^{a}	-
Acidity	$1.21{\pm}0.05^{a}$	0.4 -1.5	0.32 ± 0.01^{b}	0.25 - 0.5	0.41 ± 0.01^{b}	0.3 -0.5
Fat	0.00	0.00	0.00	0.00	0.00	0.00
Protein	$0.27{\pm}0.04^{a}$	-	0.41 ± 0.03^{a}	-	$0.50{\pm}0.01^{a}$	-
Ash	$0.54{\pm}0.03^{b}$	0.1-0.6	$0.81{\pm}0.05^{a}$	-	0.17 ± 0.01^{b}	0.1-0.3
Calcium	$0.01{\pm}0.00^{a}$	-	$0.04{\pm}0.00^{a}$	-	0.07 ± 0.01^{a}	-
Phosphorous	$0.03{\pm}0.00^{a}$	-	$0.07{\pm}0.00^{a}$	-	$0.08{\pm}0.00^{a}$	-
Brix	19.54±3.01 ^a	Min 12	12.40±3.61 ^b	Min 13.5	12.70±1.90 ^b	-
Total sugar	8.03 ± 0.89^{a}	-	$9.42{\pm}1.60^{a}$	Min 11	$7.31{\pm}1.53^{a}$	Min 12
Sugar	5.91±0.23 ^b	Min 3.5	$7.92{\pm}0.15^{a}$	-	6.22 ± 0.26^{b}	Min 6
redaction						

Properties (%)	orange	orange	grape	grape	Apple	Apple
		standard		standard		standard
Total solid	13.5±0.91 ^b	Min 10.5	$17.00{\pm}1.07^{a}$	Min 15.5	14.90 ± 3.70^{b}	Min 12.5
pН	3.18 ± 0.01^{a}	2.8 -4	3.19 ± 0.03^{a}	2.8 - 3.8	3.81 ± 0.00^{a}	3.2-4.2
Refract index	1.3615 ± 0.01^{a}	-	1.355 ± 0.01^{a}	-	$1.360{\pm}0.02^{a}$	-
Acidity	0.61 ± 0.03^{b}	Min 0.5	0.65 ± 0.03^{b}	0.35 - 1.2	0.41 ± 0.01^{b}	0.2 - 0.5
Fat	0.00	0.00	0.00	0.00	0.00	0.00
Protein	$0.26{\pm}0.02^{a}$	-	$0.23{\pm}0.01^{b}$	-	$0.32{\pm}0.05^{b}$	-
Ash	0.33 ± 0.01^{b}	0.2-0.5	0.21 ± 0.03^{b}	0.2-0.3	0.31 ± 0.02^{b}	0.2 -0.35
Calcium	$0.03{\pm}0.00^{a}$	-	$0.02{\pm}0.00^{a}$	-	$0.04{\pm}0.00^{a}$	-
Phosphorous	0.06 ± 0.00^{a}	-	$0.05{\pm}0.00^{a}$	-	$0.05{\pm}0.01^{a}$	-
Brix	18.71 ± 3.12^{a}	Min 10	14.5 ± 3.05^{b}	-	$19.01{\pm}1.09^{a}$	Min 11
Total sugar	7.12 ± 0.99^{a}	-	8.17 ± 2.64^{a}	-	$9.52{\pm}2.61^{a}$	-
Sugar redaction	5.72±0.71 ^b	Min 3.5	6.80 ± 1.01^{b}	Min 13	8.11 ± 0.87^{ab}	Min 8

Table 2: physic- chemical properties of fruit beverage