خلاصه مقالات سخنراني ويوستر



نشتی کی وری کر تحقیقات بیکاشت پاروری و گاپاروری تازه های علمی باروری و ناباروری

تهران، مرکز همایش های بین المللی دانشگاه شهید بهشتی

۲۷-۲۷ آذرماه ۱۳۹۴

# Evaluation of expression level of *BRDT* gene in testis tissues of infertile men referred to ROYAN Institute

Kohandani, F.<sup>1,2</sup>; Moshtaghioun, S.M.<sup>1</sup>; and Shahhoseini, M.<sup>2</sup>

1. Biology department, Faculty of sciences, Yazd university, Iran.

2. Department of Genetics, Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran.

### Introduction:

Spermiogenesis stage of spermatogenesis is including to many epigenetics events leading to sperm chromatin condensation. Through this condensation sperm chromatin undergoes drastic changes from nucleo-histone structure to nucleo-protamin structure, many factors are involved in this epigenetics process such as BRDT. In the post-meiotic phase of spermatogenesis, BRDT protein binds to hyperacetylated histones and aids to their general removal from DNA and their final replacement by protamine.

### Aim:

According to the role of sperm chromatin condensation in fertility and the important role of *BRDT* gene in this condensation, the expression analysis of *BRDT* gene in testis tissues of infertile men was aimed in this study.

### Material and methods:

The study included total of 30 infertile men who referred to ROYAN Institute. This samples distributed into 3 groups: obstractive azoospermia (positive control), complete maturation arrest at second spermatocyte/spermatid level and Sertoli cell syndrome (negative control). Samples of testicular tissues extracted by TESE/microTESE procedures performed in an attempt to obtain sperm, also for this respect, consent was obtained from participants. Quantitative expression data were obtained by real-time PCR technique using specific primer pair for *BRDT* gene.

### **Results:**

Our study showed *BRDT* gene expression level was significantly higher in obstractive azoospermia group compared to complete maturation arrest and Sertoli cell only syndrome groups ( $p \le 0.05$ ).

# Conclusion:

The finding in this study declared an association between low level of *BRDT* gene expression and spermatogenesis failure and male infertility.

# Keywords:

Spermatogenesis, Epigenetics, BRDT gene.