

۱

۱

۱

Comparative mosquito repellent efficacy of alcoholic extracts and essential oils of different plants against *Anopheles Stephensi*

۱

Mohammadi Ali ^a, Ghalandari Ruhollah ^a, Alizadeh Ghorban Ali ^b, Batatshoostari Ali ^c, Khatami, Mohammad Reza ^a, ۱ BaratShoostari Mohammad ^a *

^a. Biotechnology Research Center, Science and Technology Institute, Tehran, Iran.

^b. Tehran Payame-Noor University, Department of Agricultural Biotechnology, Tehran, Iran.

c. ۱ Shahrekord, Islamic Azad University, Department of Microbiology, Shahrekord, Iran.

۱

Corresponding Author;

Mohammad Barat Shoostari

Biotechnology Research Center, Science and Technology Institute, Tehran, Iran.

e-mail: behzadshoostary@gmail.com

Phone numbers: +۹۸۹۱۲۳۱۴۶۹۷۰ - +۹۸(۲۶۳)۲۳۱۱۰۸۴

۱

۱

Abstract:

Mosquitoes control and personal protection from mosquito's bites is one of the serious ways for preventing of contagious diseases distribution. Mosquitoes in addition to the local symptoms (itching, redness and irritation) can cause transmission of fatal and dangerous disease especially in tropical areas. In recent years interest in plant-based products has been revived because of the development of resistance,

cross-resistance and possible toxicity hazards associated with synthetic insecticides and their rising cost. Various plant-based products as herbal repellents are safe and biodegradable alternatives to synthetic chemicals for use against mosquitoes.

In the present study, essential oils and extracts of six plants "*Melissa officinalis*, *Rosmarinus officinalis*, *Lavandula officinalis*, *Citrus limonum*, *Eucalyptus globulus* and *Ocimum basilicum*" were evaluated compared against mosquitoes of *Anopheles Stephensi*. Finally, repellent properties of essential oils and extracts as experimental groups and of *N, N-diethyl γ -methylbenzamide (DEET)* as a positive control group were compared. We used Duncan's multiple range tests to determine the significant differences at 1% level between the experimental group and the control group. Results of statistical analysis showed significant differences between the extracts and essential oils. Essential oils indicated more effectiveness rather than extracts.

Key Words: Malaria, *Anopheles stephensi*, Insect Repellent, Essential oil, Plant extract

Introduction

Problems with chemical insecticides and possible effect of essential oils attracted the attention of researcher. In the recent years, long term application of chemical substances for controlling, repelling and killing of hazardous insects, make serious anxieties for environment and human health (Nerioa, 2010; Yang, 2002). Therefore uses of environment friendly and biodegradable natural insecticides of plant origin have received renewed attention as agents for mosquitoes control (Nerioa, 2010). The years before 1940 volatile substances such as Citronella oil, Clove seed oil and Camphor were the common insect repelling substances. In addition some plants such as *Marquise*, *eucalyptus*, *fennel*, *oregano*, *pepper*, *wormwood* plant and tea tree are known to show such properties (Wilkinson, 1996; Sima Nasri, 2012). Natural Iranian flour consists of many herbs which are traditionally used as repelling insects especially against *malaria* vectors. According to WHO report eastern Mediterranean countries like Iran are the source of contagious disease by insects specially mosquitoes species (WHO, 2010; Jinous Asgarpanah, 2012). So far 64 species of mosquitoes have been reported from Iran, which included 28 species of *Anopheles*, 3 species of *Aedes* and 19 species of *Culex* and 14 species are from other genus (Azari Hamidian, 2007). Among all the insect vectors of human disease, *Anopheles* is the most popular and worst species of mosquitoes (Collins, 1990; Curtis, 1994).

Among the species, *Anopheles stephensi* mosquito is the most important *malaria* vector especially in Iran. This mosquito in India and some of West Asian countries are

annually the reason of 40-50 percent of malaria disease. Globally, *malaria* kills 3 million people each year, including 1 child every 30 seconds (Shell, 1997). The search for effective vaccines against malaria is still in progress. Individual protecting actions, comprising repellents are extensively applied to put off the transmission of arthropod borne diseases by decreasing access between persons and vectors. (Coleman, 1993; Walker, 1996).

1

World War II and the needs of military groups in the tropical lands promoted the scientists to extensive research for discovery and preparation of insect repellent drugs (Wilkinson, 1996). Undoubtedly Iranian military forces face to such disease by the insects. Thus, more than ten thousands chemical compounds have been tested for the insect repelling effects. Nevertheless long term using of chemical insecticides has side effects on health. To avoid the adverse effect, researches on repellents that are derived from plant are promising in that they are effective, safe to users, and also inexpensive (Abu-Qare, 2001; Fradin, 1998), (Barnard, 2004). 1

For this reason and achieving better results we select the plants that can be grown in Iran and are easily propagated. An enormous amount of plant products have been stated to have mosquito larvicidal and/or repellent action against mature mosquitoes.

In the previous studies 10 medicinal herbs were used as repelling and killing of ova, larva and mature insects of *Anopheles stephensi*, *Aedes Aegypti* and *Culex quinquefasciatus*. Finally ginger and rosemary essential oils were introduced respectively as a killer and repellent of three mentioned mosquitoes. Also repellent effects of either essential oil and extract of lemon and Melissa against *Anopheles stephensi* were examined and the results showed better effect of essential oil rather than extract, but differences were not significant (Veena, 2000, Vatandoost, 2004).

1

In another study, the essential oil of five different plants leaves and their repellent effects was investigated against the *Anopheles* mosquito. Other scientists also were introduced essential oils as repelling of the malaria-carrying insect (Oshaghi, 2003) (Rajkumar, 2007).

Essential oils repellency of aromatic plants that grow in Argentina against *Aedes aegypti* have also evaluated: *Acantholippia seriphioides*, *Achyrocline satureioides*, *Aloysia citriodora*, *Anemia tomentosa*, *Baccharis spartioides*, *Chenopodium ambrosioides*, *Eucalyptus saligna*, *Hyptis mutabilis*, *Minthostachys mollis*, *Rosmarinus officinalis*, *Tagetes minuta* and *Tagetes pusilla*. Most essential oils were effective (Gillij, 2008; Muhammad Alam, 2012).

In recent years, several plants extracts including neem (*Azadirachta indica* A.Juss), Citronella grass (*Cymbopogon nardus* Rendle), basil (*Ocimum basilicum* L., *Ocimum gratissimum* L., *Ocimum americanum* L.), clove (*Syzygium aromaticum* L.), prickly straggler (*Solanum trilobatum* L.), musk basil (*Moschosma polystachyum* L.) and

thyme (*Thymus vulgaris* L.) have been studied as possible mosquito repellents. (Gillij, ۲۰۰۷).

Consequently, the aim of this study is to compare the repellent activity of plant essential oils and plant extract of six plants against the *Anopheles stephensi*.

۱

Materials and Methods

۱

۱ - Plant selection

۱

In this study plants according to ancient data bases for insect repelling were selected. Also tried to choose plants that are easily propagated and using of them be safe for human and doesn't induce toxicity in person.

۱

The below table is followed by six plants were tested.

name Common	used for Parts	Family	name Plant	No
Oils Extracts and Essential				
Lavender	Flower	Lamiaceae	officinalis Lavandula	۱
Melissa	Leaf	Labiatae	officinalis Melissa	۲
Rosemary	Leaf	Lamiaceae	officinalis Rosmarinus	۳
Lemon	Peel	Rutaceae	limonum Citrus	۴
Eucalyptus	Leaf	Myrtaceae	globolus Eucalyptus	۵
Basil	Herb	Lamiaceae	basilicum Ocimum	۶

۱

The plants from the farm of Karaj Agriculture Faculty (Tehran University) were prepared. Leaves and spire of the plants which were contain the effective substances, were selected and in a proper condition with suitable air circulation and far from direct sun light were dried. Dried herbal parts were milled. Then from powder by the maceration method and use of ۷۰٪ ethanol, alcoholic extract were prepared. In this method ۱۰۰g powder in one liter of ethanol as a solvent macerated then by vacuum machine and in temperature of ۴۰°C were concentrated.

For essential oil preparing, for each ۱۰۰ grams of herbal powders, one liter double distilled water was added and essential oil extraction was done in Clevenger machine (Tyler , ۱۹۸۸). Then repellent effects of both herbal extracts and essential oils separately on Anopheles mosquitoes were investigated.(Buescher, ۱۹۸۵ تا Klun, ۲۰۰۰).

To prepare different concentrations, the products were further diluted using alcohol as diluents. Extract and essential oil solutions were formulated on a volume- volume basis at a concentration of 3% and 1%, respectively. The compounds were applied as 4-ml aliquots of ethanol solution and were spread evenly over the animal skin as explained previously. (Buescher, 1980 and Klun, 2000).

۱

۲. Insect Selection

The mosquitoes of *Anopheles stephensi* obtained from a well-established laboratory colony from school of Public Health & Institute of Health Research, Tehran University of Medical Sciences. Mosquitoes were reared and maintained at $27\pm 1^{\circ}\text{C}$ and $80\pm 1\%$ Relative Humidity(RH) under a 12:12(L:D) photoperiod. Larvae were reared on a diet of floating catfish food. Female mosquitoes are only the *malaria* vectors, so they selected. The adults were maintained in screen cages on 10% sucrose solution but 24 hours before experiments the sucrose solution was removed from cages. Repellency was tested against 3-day-old, blood-starved mosquitoes, and for each test 20 mosquitoes were used. (Barat hooshtari, 2012 and Norashiqin, 2008).

۱

Animal testing

The extracts and essential oils were tested on animal guinea pigs. The animals were laboratory reared albino males with average 400-450 g weight. A 4-5 cm of animal abdomen hairs was removed then washed and cleaned by ethanol. Treatments were 4 ml of either the extracts containing 0.12 and 0.04 g of active ingredients of extracts and oils respectively, and 0.4 ml DEET (*N, N-diethyl-3-methylbenzamide*) in the same way, were used as a positive control. After treatment, the animal was bound on top of the cage in which the treated position was exposed to mosquitoes for thirty minutes. Each test was repeated three times replacing new mosquitoes and new animal, and number of bites through the fabrics was recorded. Animal experimentation was approved by the Animal Research Committee of Tehran University.

۱

Statistical analysis

To compare the repellent efficacy of the compounds, we used Duncan's multiple range tests to determine the significant differences at 1% level between the experimental group and the control group.

۱

Results

Results of laboratory study on animals comparing extracts and oils of *Melissa officinalis*, *Rosmarinus officinalis*, *Lavandula officinalis*, *Citrus limonum*, *eucalyptus globulus*, *Ocimum basilicum* and DEET against *A. Stephensi* are presented in table ۱. Results showed that oils were significantly more effective than extracts. *Lavandula officinalis* and *eucalyptus globulus* oils have most effectiveness. In this study there was no significant differences between oil and extract of *Lavandula* (۹۷,۱۶% and ۹۶,۵۷% respectively) and *Eucalyptus* (۹۷,۱۵% and ۹۰,۰۶% respectively) that could be indicate the better effectiveness of these two herbs in repelling of *Anopheles Stephensi*. There are significant differences in the effectiveness rate of oils and extracts of *Melissa officinalis*, *Rosmarinus officinalis*, *Ocimum basilicum* and especially *Citrus limonum*, which repelling effect of the extracts were less than oils (Table ۲). In this study comparing repellent effect of oils rather than chemical control (*DEET*) was acceptable but comparing of *DEET* with extracts was not significant.

۱

۱

۱

۱

۱

۱

۱

۱

۱

۱

۱

۱

۱

۱

Table ۱. Relative repellent effectiveness of ۳% extracts and ۱% oils of *Lavandula officinalis*, *Melissa officinalis*, *Rosmarinus officinalis*, *Citrus limonum*, *Eucalyptus globulus*, *Ocimum basilicum* laboratory mosquitoes of *Anopheles stephensi* on guinea-pigs in the laboratory. Efficacy of each experimental substance was repeated three times. Highest efficacy were supposed for *Lavender oil* and lowest efficacy for *Ethanol*.

(%)SE	Mean	III Replicate	Replicate II	Replicate I	Repellent
۲,۷۵±۱۹۷,۱۶	a ۹۷,۱۶۶۷	۹۴,۵	۹۷	۱۰۰	oil Lavender
۷,۷۸±۱۹۲,۶۶	a ۹۲,۶۶۶۷	۱۰۰	۸۴,۵۰	۹۳,۵	oil Melissa
۶,۶۵±۱۹۱,۶۳	a ۹۱,۶۳۳۳	۹۷,۵	۹۳	۸۴,۴۰	oil Rosemary
۳,۰۰±۱۹۲,۶۳	a ۹۲,۶۳۳۳	۹۲	۹۰	۹۵,۹	oil Lemon
۲,۴۸±۱۹۷,۱۵	a ۹۷,۱۵۰۰	۱۰۰	۹۵,۴۵	۹۶	oil Eucalyptus
۱,۵۵±۱۹۴,۶۶	a ۹۴,۶۶۶۷	۹۴,۲	۹۳,۴	۹۶,۴	Basil oil
۳,۰۰±۱۹۷,۰۰	a ۹۷,۰۰۰۰	۹۴	۱۰۰	۹۷	(DEET(standard
۲,۹۶±۱۹۶,۵۷	a ۹۶,۵۷۶۷	۹۴,۷۳	۱۰۰	۹۵	extract Lavender
۶,۲۱±۱۷۶,۳۰	a ۹۰,۰۶۶۷	۷۴,۲	۸۳,۳	۷۱,۴۲	extract Melissa
۶,۲۸±۱۷۲,۳۲	b ۷۶,۳۰۶۷	۶۶,۸	۷۱	۷۹,۱۶	extract Rosemary
۱۸,۰۱±۱۵۰,۱۲	b ۷۲,۳۲۰۰	۶۲,۱۵	۵۸,۸	۲۹,۴۱	extract Lemon
۵,۲±۱۹۰,۰۶	b ۷۳,۹۳۳۳	۸۴	۹۲,۵	۹۳,۷	extract Eucalyptus
۸,۲±۱۷۳,۹۳	c ۵۰,۱۲۰۰	۶۶	۸۲,۵	۷۳,۳	extract Basil
.	(Ethanol(control

İ Average that have the same letters are not statistically significant at ۵%.

ظEfficacy of each experimental substances was repeated three times and finally we calculate the mean.

İ

İ

İ

İ

Table ۲. Influence of variance analysis of extracts and essential oils of six herbs on the repellent rate of *Anopheles stephensi* in guinea pigs.

F _C	۱%F	۵%F	df	Square Sum of	Squares Mean	of Variant Source
**۱۲,۰۹	۲,۹۶	۲,۱۵	۱۲	۷۲۱۹,۲۳	۶۰۱,۶۰	(Groups between)Treatment
			۲۶	۱۲۹۳,۵۶۷	۴۹,۷۵	(Groups Within) Error

**Significant at ۱% probability

İ

İ

Discussion

Present study shows herbal essential oils have better repellent efficacy rather than herbal extracts. Hence essential oils could be also used as a better and safer substitution of chemical repellent substances such as *DEET* (Barat shooshtari,۲۰۱۲). previous studies regarding the extracts and essential oils of *Melissa*, *Rosemary*,

Lavender, *lime*, and *ginger* that have done by others support our study and they have also proposed the essential oils as a suitable substitution of chemical repellent (Oshaghi, 2003; Rajkumar, 2007; Barat shooshtari, 2012; Kweka, 2009).

In the animal experiments, *Lavender* and *Eucalyptus* oils rather than other oils had a better repellent effectiveness, 97.16% and 97.15% respectively, against *anopheles*. (Gillij, 2007). Therefore they could be recommended as a safe and suitable substitution of chemical repellent. In this work, we have tested the repellents against only one species and do not know if these compounds are protective against other mosquito species or medically important insects. By more clinical trial may introduce the essential oils in the insect's repellent herbal cream formulation. Different factors may interfere in insect's repellent efficacy that the main factor could be effective substances of essential oils and extracts. Therefore analysis of different fractions of herbal oils and extracts and its effect on the insects is recommended. Most of these essential oils are highly volatile and this contributes to their poor longevity as mosquito repellents. However, this problem can be addressed by using fixatives or careful formulation to improve their longevity. For example, oils from turmeric and hairy basil with addition of 0% vanillin repelled 3 species of mosquitoes under cage conditions for a period of 6-8 hours depending on the mosquito species (Tawatsin, 2001). The exception to this is *para-menthane* 3, 4 *diol*, which has a lower vapour pressure than volatile *monoterpenes* found in most plant oils (Barasa, 2002) and provides very high protection from a broad range of insect vectors over several hours (Carroll, 2006).

The plants can be used alone or combined for effective protection against mosquitoes. They can also be used for control of mosquito breeding. (Barnard, 2004; Trongtokit, 2005). They also offer safer alternative to synthetic chemicals and can be obtained by individuals and communities easily at a very low cost. However toxicity tests of the active plants need to be done to ascertain their safety in administration. (Robert, 1991; Rutledge, 1978).

p

p

p

Conclusion

The identification of these potential repellent plants from the local flora will generate local employment and stimulate local efforts to enhance public health. However, pilot studies are indicated to evaluate the epidemiological impact and cost-effectiveness of the natural oils, which are reported to be effective in mosquito control or provide protection against mosquito bites. Further investigations are needed to elucidate the six essential oils against a wide range of mosquito species and also to identify active

compounds responsible for repellent activity and to be utilized if possible, in preparing a commercial product/formulation to be used as insecticidal.

۱

References:

۱ Abu-Qare, A. and Abou-Donia MB. ۲۰۰۱. Development of high performance liquid chromatographic method for the quantification of chlorpyrifos, pyridostigmine bromide, N, N-diethyl toluamide and their metabolites in rat plasma and urine. J Chromatography B, ۷۵۴, ۵۳۳-۵۳۸.

۱ Azari Hamidian, S., ۲۰۰۷. Checklist of Iranian mosquitoes (Diptera: Culicidae). J of Vec Ecol. ۲۲(۲): ۲۳۵-۲۴۲.

۱ Barasa SS, Ndiege IO, Lwande W, Hassanali A. ۲۰۰۲. Repellent activities of stereoisomers of p-menthane-۳,۸-diols against Anopheles gambiae (Diptera: Culicidae). J Med Entomol ۳۹:۷۳۶-۷۴۱.

۱ Barat shooshtari, M. and Galandari, R. ۲۰۱۲. Comparative study on Repellent Effect of Extracts and Essential Oils of Melissa officinalis, Rosmarinus officinalis and Lavandula angustifolia Against Main Malaria Vector, Anopheles stephensi (Diptera: Culicidae). Iranian journal of Medicinal and Aromatic plant. Vol. ۲۷, No. (۴): ۶۰۶-۶۱۳.

۱ Barnard, DR. and Dexue, RD., ۲۰۰۴. Laboratory evaluation of mosquito repellents against Aedes albopictus, Culex nigripalpus, and Ochlerotatus triseriatus (Diptera: Culicidae). J Med Entomol ۴۱:۷۲۶-۷۳۰.

۱ Bohbot, J.D., et al., *Multiple activities of insect repellents on odorant receptors in mosquitoes*. Med Vet Entomol, ۲۰۱۱. ۲۵(۴): p. ۴۳۶-۴۴۴.

۱

۱ Buescher, MD. Rutledge, LC. Writz, RA. Nelson, JH., ۱۹۸۵. Laboratory repellent tests against Rhodnius prolixus. J Med Ent, ۲۲: ۴۹-۵۳.

۱ Carroll SP, Loye J. ۲۰۰۶. PMD, a registered botanical mosquito repellent with deet-like efficacy. J Am Mosq Control Assoc, ۲۲:۵۰۷-۵۱۴.

۱ Coleman, R.E. Robert, L.L. Roberts, L.W. Glass, J.A. Seeley, D.C. Laughinghouse, A. Perkins, P.V., and Wirtz, R.A., ۱۹۹۳. Laboratory evaluation of repellents against four anopheline mosquitoes (Diptera: Culicidae) and two Phebotomine sand flies (Diptera: Psychodidae). Journal of Medical Entomology ۳۰: ۴۹۹-۵۰۲.

1 Collins, F.H. and Paskewitz, S.M., 1990. Malaria: Current and future prospects for control. Annual Review of Entomology. 40: 190-219.

1 Curtis, C.F., 1994. Should DDT continue to be recommended for malaria vector control?. Medical and Veterinary Entomology. 8: 107-112.

1 Fradin, M., 1998. Mosquitoes and mosquito repellent, Annals of Internal Medicine. Mosquito Repellent with DEET, <http://www.dphhs.state.mt.us/news/west-nile-smosquito-repellant-deet.htm>. 128(11), 931-40.

1 Gillij, Y.G. Gleiser, R.M. Zygadlo, J.A. Mosquito repellent activity of essential oils of aromatic plants growing in Argentina. [Bioresource Technology](#). [Volume 99, Issue 7](#), May 2008, Pages 2007-2010.

Jinous Asgarpanah and Fereshteh Ramezanloo. Chemistry, pharmacology and medicinal properties of Peganum harmala L. African Journal of Pharmacy and Pharmacology Vol. 6(22), pp. 1073-1080, 10 June, 2012.

1 Klun, JA, and Debboun, M., 2000. A new model for quantitative evaluation of repellents efficacy using human subjects. J Med Ent, 37: 177- 81.

1 Kweka, E.J, Mwangonde, B. J. Mahande, M.J. Mosha, F.W. Kimaro, E. E. S. Msangi, L. Lyaruu and A. M. Mahande., 2009. Evaluation of Ocimum suave essential oil against anthropophilic malaria vector Anopheles gambiae s.s after ten years of storage. Tanzania Journal of Health Research, Vol. 11, No. 2.

Muhammad Alam, Ghiasuddin, Anwar Sadat, Naveed muhammd, Ashfaq Ahmad Khan and Bina S. Siddiqui. Evaluation of Viburnum grandiflorum for its in-vitro pharmacological screening. African Journal of Pharmacy and Pharmacology Vol. 6(22), pp. 1606-1610, 10 June, 2012.

1 Nerioa, LS.1 Olivero-Verbel J. and Stashenko E., 2010. Repellent activity of essential oils: A review. [Bioresource Technology](#) [Volume 111, Issue 1](#), January. Pages 372-378.

1 Norashiqin, M. and1 Sallehudin, S., 2008. The Repellent Activity of Piper aduncum Linn (Family: Piperaceae) Essential Oil against Aedes aegypti Using Human Volunteers. Available online at www.ptat.thaigov.net. J Trop Med Parasitol. 31:63-9.

1 Oshaghi, MA. Ghalandari, R. Vatandoost, H. Shayeghi, M. Kamali-nejad, M. Tourabi-Khaledi, H. Abolhassani, M. and Hashemzadeh, M., 2003. Repellent Effect of Extracts and Essential Oils of Citrus limon (Rutaceae) and Melissa officinalis (Labiatae) Against Main Malaria Vector, Anopheles stephensi (Diptera: Culicidae). Iranian J Pub I Health, Vol. 32, No. 4, pp.47-52.

1 Rajkumar, S. and Jebanesan, A., 2007. Repellent activity of selected plant essential oils against the malarial fever mosquito *Anopheles stephensi*. *Tropical Biomedicine* 22(2): 71-75.

1 Robert, LL. Hallan, JA. Seeley, DC. Roberts, LW. and Wirtz, RA., 1991. Comparative sensitivities of four *Anopheles* (Diptera: Culicidae) to five repellents. *J Med Ent*, 28: 417-20.

1 Rutledge, LC. Moussa, MA. Lowe, CA. and Sofield, RK., 1978. Comparative sensitivity of mosquito species and strains to the repellent diethyl toluamide. *J Med Ent*, 15: 536-41.

1 Shell, E.R., 1997. Resurgence of a deadly diseases. *Atlantic Monthly*. 40-60.

Sima Nasri, Mahdieh Anoush and Narges Khatami. Evaluation of analgesic and anti-inflammatory effects of fresh onion juice in experimental animals. *African Journal of Pharmacy and Pharmacology* Vol. 6(23), pp. 1679-1684, 22 June, 2012.

1

1 Tawatsin A, Wratten SD, Scott RR, Thavara U, Techadamrongsin Y: 2001. Repellency of volatile oils from plants against three mosquito vectors. *J Vector Ecol*, 26:76-82.

1 Trongtokit, Y. Rongsriyam, Y. Komalamisra, N. and Apiwathnasorn, C., 2005. Comparative repellency of 38 essential oils against mosquito bites. *Phytother Res*. 19:303-9.

1 Tyler, E. Brady, LR. and Robbers, JE., 1988. Volatiles oils, In: *Pharmacognosy*. Eds, Lea & Febiger. 9th Ed. Philadelphia, pp. 100-107.

1 Vasudevan, P. Tandon, M. Pathak, N. Nuennerich, P. Mueller, F. and Mele, A., 1997. Fluid CO₂ extraction and hydro distillation of certain biocidal essential oils and their constituents. *J Sci Ind Res*. 56:662-72.

1 Vatandoost, H. and V.M. Vaziri, *Larvicidal activity of a neem tree extract (Neemarin) against mosquito larvae in the Islamic Republic of Iran*. *East Mediterr Health J*, 2004. 10(4-5): p. 573-81.

1

1 Veena Prajapati, A.K. Tripathi, K.K. Aggarwal and S.P.S., 2005. Khanuja. Insecticidal, repellent and oviposition-deterrent activity of selected essential oils against *Anopheles stephensi*, *Aedes aegypti* and *Culex quinquefasciatus*, P.O. CIMAP, Lucknow, India [Bioresearch Technology](http://www.BioresearchTechnology.com) Central Institute of Medicinal

and Aromatic Plants, P.O. CIMAP, Lucknow, India- [Bioresearch Technology](#).
Pages ۱۷۴۹-۱۷۵۷, Volume

۱ Walker, T.W. Robert, L.L. Copeland, R.A. Githeko, A.K. Wirtz, R.A. Githure, J.I. and Klein, T.A., ۱۹۹۶. Field evaluation of arthropod repellents, DEET and a piperidine compound, AB-۳۷۲۲۰, against *Anopheles funestus* and *Anopheles arabiensis* in western Kenya. *Journal of American Mosquito Control Association* ۱۲: ۱۷۲-۱۷۶.

۱ Wilkinson, J.B., ۱۹۹۶. Insect repellents. In: Wilkinson J.B.; Moor R.J. *Harrys cosmeticology: From Longman Scientific & Technical*. Singapore: ۷th ed. ۲۰۷-۸.

۱ World Health Organization, ۲۰۰۴. Integrated vector managment. Cairo. WHO Regional Officer for Eastern Mediterranean.

۱ Yang, YC. Lee, SG. Lee, HK. Kim, MK. Lee, SH. Lee, HS., ۲۰۰۲. A piperidine amide extracted from *Piper longum* L. fruit shows activity against *Aedes Aegypti* mosquito larvae. *J Agric Food Chem*. ۵۰: ۳۷۶۵-۷.

۱

۱

۱

۱

۱