

Weak Almost Periodicity and Uniform Continuity on a Locally Compact Quantum Group

R. FAAL and H. R. EBRAHIMI VISHKI*

Abstract

First we present some useful characterizations for the function spaces $AP(\mathbb{G})$, $WAP(\mathbb{G})$ and $LUC(\mathbb{G})$ on a locally compact quantum group \mathbb{G} and then we study some conditions under which $AP(\mathbb{G})$ and $WAP(\mathbb{G})$ are C^* -algebras. We also investigate the equality $LUC(\mathbb{G}) = WAP(\mathbb{G})$.

2010 *Mathematics subject classification*: Primary 47L10, Secondary 46H25.

Keywords and phrases: Locally compact quantum group, coamenable, (weak) almost periodicity, left uniform continuity.

1. Introduction

Let G be a locally compact group. It is known that the function spaces $AP(G)$, $WAP(G)$ and $LUC(G)$ are C^* -subalgebras of $C_b(G)$. Moreover, they enjoy the inclusion relations $AP(G) \subseteq WAP(G) \subseteq LUC(G)$, where the equality hold under some additional conditions. For example, it has been known that $WAP(G) = LUC(G)$ if and only if G is compact. From the operator theory point of view, it is known that $WAP(G)$ can be identified with $WAP(L^1(G))$. Also there is a characterization of left uniform continuity, namely $LUC(G) = L^\infty(G) \cdot L^1(G)$. For complete information with more details on these function spaces and their inclusion relationships one may refer to [1].

We refer to [4] for the definitions and basic facts on Hopf von Neumann algebras and locally compact quantum groups. In [2] Daws showed that for an abelian Hopf von Neumann algebra (M, Γ) , $WAP(M_*) = \{x \in M : L_x : M_* \rightarrow M \text{ is weakly compact}\}$ and $AP(M_*) = \{x \in M : L_x : M_* \rightarrow M \text{ is compact}\}$ are C^* -subalgebras of M . In [3] he defined $WAP(M, \Gamma)$ for an arbitrary Hopf von Neumann algebra (M, Γ) and proved that it is the largest C^* -algebra contained in $WAP(M_*)$. However, in contrast to the situation for group, it was not known whether the equality $WAP(M_*) = WAP(M, \Gamma)$ holds. In this talk we show that for a coamenable locally compact quantum group \mathbb{G} , the spaces $WAP(L^1(\mathbb{G}))$ and $AP(L^1(\mathbb{G}))$ are C^* -algebras and that $WAP(M_*) = WAP(M, \Gamma)$. We also define the notation of completely weakly compact operator for an arbitrary

* speaker

Hopf von Neumann algebra and we give a characterization of $WAP(M, \Gamma)$ in terms of complete weak compactness of left translation operators. On the other hand Runde [5] showed that for certain locally compact quantum groups \mathbb{G} , $LUC(\mathbb{G})$ is a C^* -algebra. In this talk we also improve Runde's result.

2. Main Results

For a Hopf von Neumann algebra M we define $CWAP(M_*)$ and we show the following result.

Theorem 2.1. *Let M be a Hopf von Neumann algebra. Then*

- (i) $CWAP(M_*)$ is equal to $WAP(M, \Gamma)$, and it is the largest C^* -algebra in $WAP(M_*)$.
- (ii) If M is abelian, then $WAP(M_*) = CWAP(M_*)$, so $WAP(M_*)$ is a C^* -subalgebra of M .

We give some new characterizations for $WAP(L^1(\mathbb{G}))$, $AP(L^1(\mathbb{G}))$ and $LUC(\mathbb{G})$ from which we present the following results.

Theorem 2.2. *Let \mathbb{G} be a coamenable locally compact quantum group. Then $WAP(L^1(\mathbb{G}))$ and $AP(L^1(\mathbb{G}))$ are C^* -algebras.*

Theorem 2.3. *Let \mathbb{G} be a locally compact quantum group such that $L^1(\mathbb{G})$ is strongly Arens irregular. Then $WAP(L^1(\mathbb{G})) = LUC(\mathbb{G})$ implies that \mathbb{G} is compact.*

References

- [1] J. BERGLUND, H. JUNGHEHN AND P. MILNES, Analysis on Semigroups; Function Spaces, Compactifications, Representations, Wiley, New York, 1989.
- [2] M. DAWS, Characterising weakly almost periodic functionals on the measure algebra, *Studia Math.* **204** (2011) 213-234.
- [3] M. DAWS, Non-commutative separate continuity and weakly almost periodicity for Hopf von Neumann algebras, *J. Funct. Anal.* **269** (2015) 683-704.
- [4] J. KUSTERMANS AND S. VAES, Locally compact quantum groups in the von Neumann algebraic setting, *Math. Scand.* **92** (2003), 68-92.
- [5] V. RUNDE, Uniform continuity over locally compact quantum groups, *J. London Math. Soc.* **80** (2009) 55-71.

R. FAAL,
Department of Pure Mathematics, Ferdowsi University of Mashhad,
P. O. Box 1159, Mashhad 91775, Iran.
e-mail: faal.ramin@yahoo.com

H. R. EBRAHIMI VISHKI,
Department of Pure Mathematics and Centre of Excellence in Analysis on Algebraic Structures (CEAAS), Ferdowsi University of Mashhad,
P. O. Box 1159, Mashhad 91775, Iran.
e-mail: vishki@um.ac.ir